
LAND SUITABILITY MAP

ROBUSTA, LIBERICA AND EXCELSA COFFEE

LAND RESOURCES EVALUATION AND SUITABILITY ASSESSMENT OF STRATEGIC PRODUCTION AREAS


PROVINCE OF AGUSAN DEL NORTE





MAP INDEX

LAND RESOURCES EVALUATION AND SUITABILITY ASSESSMENT OF STRATEGIC PRODUCTION AREAS PROVINCE OF AGUSAN DEL NORTE



LAND SUITABILITY MAP FOR ROBUSTA, LIBERICA AND EXCELSA COFFEE

LAND RESOURCES EVALUATION AND SUITABILITY ASSESSMENT OF STRATEGIC PRODUCTION AREAS

AGUSAN DEL NORTE, REGION XIII

EXTENT OF SUITABILITY FOR ROBUSTA, LIBERICA AND EXCELSA COFFEE PRODUCTION BY MUNICIPALITY

	EXISTING COFFEE (Ha)				EXPANSION AREA (Ha)						CONFLICT RESOLUTION AREA (Ha)					TOTAL	
MUNICIPALITY				TOTAL EXISTING AREA (Ha)	Coconut		Shrubland, unmanaged*		Grassland, unmanaged*		Corn		Paddy rice, non-irrigated		Other crops		POTENTIAL EXPANSION AREA (Ha)
	S1	S2	S 3		S1	S2	S1	S2	S1	S2	S1	S2	S1	S2	S1	S2	AREA (IIa)
BUENAVISTA	-		-	-	7,898	103	434	130	892	80	680	-	-	-	-	-	10,216
BUTUAN CITY	-		-	-	20,582	2	249	4	200	-	7,145	-	-	-	-	-	28,181
CARMEN	-		-	-	1,839	26	-	-	362	1	141	-	-	-	-	-	2,370
CITY OF CABADBARAN	-		-	-	3,703	-	-	-	199	-	2,974	-	-	-	-	-	6,876
JABONGA	-		-	-	1,031	234	-	-	109	16	820	-	-	-	-	-	2,210
KITCHARAO	-		-	-	1,038	29	15	1	24	6	307	-	-	-	-	-	1,420
LAS NIEVES	-	1	-	-	6,345	365	450	-	319	-	1,381	7	-	-	-	-	8,868
MAGALLANES	-		-	-	815	-	-	-	-	-	183	-	-	-	-	-	997
NASIPIT	-		-	-	1,175	23	-	-	616	29	420	-	-	-	-	-	2,262
REMEDIOS T. ROMUALDEZ	-		-	-	874	-	-	-	-	-	543	-	-	-	-	-	1,417
SANTIAGO	-		-	-	1,417	-	_	-	159	-	460	-	-	_	-	_	2,036
TUBAY	-		-	-	1,996	339	_	-	13	164	445	-	-	-	-	_	2,957
TOTAL	-	-	-	1	48,714	1,121	1,148	134	2,893	296	15,498	7	-	-	-	-	69,809

Note: Delivery of robusta coffee planting materials must be started on the onset of rainy season.

<u>10</u> F2-Tc

20 T2-E3-Sh2-Rc3

30 T3-E3-Rc2

AGRONOMIC REQUIREMENT OF ROBUSTA, LIBERICA AND EXCELSA COFFEE PRODUCTION

LAND UTILIZATI TYPE	ON SUITABILITY RATING	SLOPE (%)	SOIL DEPTH (cm)	SOIL TEXTURE	SOIL DRAINAGE	SOIL REACTI (pH)		FLOODING CLASS	EROSION CLASS	ROCK OUTCROPS	ELEVATION (masl)	ANNUAL RAINFALL (mm)	CLIMATIC TYPE
Coffee	S1	<8	>100	CL, SiCL, SCL, SC, SiC, C, HC	WD,MWD	5.6 -7.2	2 high	none-slight	none-slight	none-few	<1000	2001-4500	I, III, IV
(Robusta Excelsa,	S2	8 - 30	30 - 100	FSL, L, SiL	SPD,PD	5.1 - 5. 7.3 - 7.	1 medilim	moderate	moderate	common	1000-2000	1000-2000	I, II
Liberica)	S3	>30	<30	S, LS, CSL, SL	VPD,ED	<5.0 -> 7	7.9 low	severe	severe	many	>2000	<1000 >4500	
SLOPE (%) SOIL DRAINAGE						SOIL REA	CTION (pH)		SOIL TEXTURE				
0 - 3 - level to gently sloping ED -				excessively drained		< 4.5	- extremely acid		Coarse			Fine	
3 - 8 - gently sloping to undulating			WD -	well drained		4.5 - 5.0	 very strongly acid 		S -	- sand		SC -:	sandy clay
8 - 18 - undulating to rolling		MWD -	5			5.1 - 5.5 - strongly acid			loamy sand		SiC -:	silty clay	
18 - 30 - rolling to moderately steep		SPD - somewhat poorly drained			5.6 - 6.0 - medium acid				- coarse sandy loam			clay	
30 - 50	· steep		PD -	poorly drained		6.1 - 6.5	 slightly acid 		SL -	- sandy loam		HC -	heavy clay
> 50	very steep	o VP		VPD - very poorly drained			6.6 - 7.2 - neutral						
						7.3 - 7.8	 mildly alkaline 		FSL -	fine sandy loam			
SOIL DEPTH (cm)			SURFACE IMPEDIMENT			7.9 - 8.4 - moderately alkaline			L -	· loam			
0 - 30 - very shallow			ROCK OUTCROPS			> 8.5	- strongly alkaline		SiL -	- silt loam			
30 - 50	shallow		< 10% -	none - few					CL -	- clay loam			
50 - 100	moderately deep		10 - 30% -	common					SiCL -	silty clay loam			
> 100	deep to very deep		> 30% -	many					SCL -	- sandy clay loam			

LAN	LAND LIMITATIONS DESCRIPTION AND COMBINATIONS												
ELEVA	ΓΙΟΝ		SOIL DR	AINAGE		SOIL DEPTH					EROSION		
El2 -	El2 - 1000m - 2000m			D2 - Somewhat poorly drained to poorly drained					moderately deep (30 - 10	00cm) E2	- Moderate erosion		
El3 -	El3 -> 2000m D3 - Vo			ery poorl	y drained or excessively	drained	Sh3	Sh3 - Very shallow (< 30cm)			- Severe erosion		
SLOPE	SLOPE/TOPOGRAPHY SOIL TEXTURE						ROCI	FLOC	FLOODING				
	•			Tc - Coarse texture				- Common		F2	- Moderate seasonal flooding		
								- Many	F3	- Severe seasonal flooding			
CODE	LIMITATION	CODE	LIMITATION	CODE	I INVERT A TROPI	CODE	LIBARTATION	CODE	LIMITATION	CODE	LANDUCE		
CODE	LIMITATION	CODE	LIMITATION	CODE	LIMITATION	CODE	LIMITATION	CODE	LIMITATION	CODE	LANDUSE		
1	E2	11	F3-D2	21	T2-El2	31	T3-E3-Sh2-Rc3	41	T3	4	Corn		
2	E2-Rc3	12	Rc2	22	T2-El2-E3	32	T3-E3-Sh3-Rc2	42	T3-E3	81	Coffee		
3	E2-Sh2-Rc2	13	Sh2	23	T2-El2-E3-Rc2	33	T3-E3-Sh3-Rc3	43	T3-E3-Rc3	82	Cacao		
4	E3-Rc3	14	Sh2-Rc2	24	T2-El2-E3-Rc3	34	T3-El2	44	T3-E3-Sh3-Rc3	116	Coconut		
5	E3-Sh2-Rc3	15	T2	25	T2-El2-E3-Sh2-Rc2	35	T3-E12-E3	45	T3-E12	126	Grassland		
6	El2-E3-Rc3	16	T2-E3	26	T2-F2-D2	36	T3-El2-E3-Rc2	46	T3-El2-E3	130	Bare areas, unmanaged		
7	El2-Rc2	17	T2-E3-Rc2	27	T2-F3-D2	37	T3-El2-E3-Sh2-Rc3	47	T3-El2-E3-Rc3	134	Shrubs, unmanaged		
8	El2-Sh2-Rc2	18	T2-E3-Rc3	28	T3	38	T3-El2-E3-Sh3-Rc2	48	T3-El2-E3-Sh3-Rc3				
9	F2-D2	19	T2-E3-Sh2-Rc2	29	T3-E3	39	T3-El2-E3-Sh3-Rc3	49	T3-El3				
1	1	1		1	1	1	1	1	I I				

40 T3-F3-D2

SUITABILITY CLASSES:

Highly Suitable (S1) Land having no significant limitation to sustained application of a given use, or only minor limitations that will not significantly reduce productivity or benefits and will not raise inputs above an acceptable level.

Marginally Suitable (S3) Land having limitations which in aggregate are severe for sustained application of a given use and will so reduce productivity or benefits, or increase required inputs, that this expenditure will be only marginally justified.

Moderately Suitable (S2) Land having limitation which in aggregate are moderately severe for sustained application of a given use; the limitation will reduce productivity or benefits and increase required inputs to the extent that the overall advantage to be gained from the use, although still attractive, will be appreciably inferior to that expected on class S1 land.

Not Suitable / Not Relevant Land having limitations which may be surmountable in time but which cannot be corrected with existing knowledge at currently acceptable cost; the limitations are so severe as to preclude successful sustained use of the land in the given manner. Existing forest, shrubland greater than 18% slope, irrigated paddy rice and miscellaneous land types such as built up areas, roads, etc are considered as not relevant.

CLIMATE TYPE


TYPE I: Two pronouced season, dry from November to April and wet during the rest of the year. Maximum rain period is from June to September

TYPE II: No dry season with a very pronounced maximum rain period from December to February. There is not a single dry month. Maximum monthly rainfall occurs during the period from March to May.

TYPE III: No very pronounced maximum rain period, with a dry season lasting only from one to three months, either during the period from December to February or from March to May. This type resembles Type I since it has a short dry season.

TYPE IV: Rainfall is more or less evenly distributed throughout the year. This type resembles Type II since it has no dry

Western part of Agusan Del Norte is classified as climatic Type IV and North Eastern part is climatic Type II.



Source: PAGASA 2018, Climatological Normals (Rainfall), Philippine Atmospheric, Geophysical and Astronomical Services Administration (PAGASA), accessed 27 July 2018, https://www1.pagasa.dost.gov.ph/index.php/climate/climatological-normals.

^{*}establishment of shade trees prior to planting of robusta coffee.

