



Department of Agriculture
BUREAU OF SOILS AND WATER MANAGEMENT
"NATIONAL SOIL SAMPLING AND TESTING FOR FERTILITY AND CROP SUITABILITY ASSESSMENT"
FERTILIZER-GUIDE FOR RICE
PROVINCE OF MISAMIS ORIENTAL

| Barangay | Farmer Representative | Fertilizer-Guide Group | | Additional 10-20 bags/ha Organic Fertilizer is recommended for all options | | | | | | | | | | | | | | | | | | | | |
|----------------------|----------------------------------|------------------------|-------------|----------------------------------------------------------------------------|----|----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|----|------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|----------------------|----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------|----|----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--|--|------------|--|--|
| | | | | INBRED RICE | | | | | | | | | HYBRID RICE | | | | | | | | | | | |
| | | Number | Description | WET SEASON | | | Nutrient Requirement | | | DRY SEASON | | | Nutrient Requirement | | | WET SEASON | | | Nutrient Requirement | | | DRY SEASON | | |
| | | | | | | | | | | | | | | | | | | | | | | | | |
| Municipality of OPOl | | | | | | | | | | | | | | | | | | | | | | | | |
| Taboc | Apolinar Daasin, Alberto Saludes | 080 | M-L-H-H | 90 | 7 | 7 | Option 1: (1st Application): 1.00 (14-14-14); 1.75 (46-0-0); (2nd Application): 1.00 (46-0-0); (3rd Application): 1.00 (46-0-0); Option 2: (1st Application): 0.75 (16-20-0); (2nd Application): 2.00 (46-0-0); (3rd Application): 1.75 (46-0-0); 0.25 (0-0-60); Option 3: (1st Application): 2.00 (46-0-0); 1.00 (0-18-0); (2nd Application): 1.00 (46-0-0); (3rd Application): 1.00 (46-0-0); 0.25 (0-0-60); | 100 | 7 | 7 | Option 1: (1st Application): 1.00 (14-14-14); 2.00 (46-0-0); (2nd Application): 1.25 (46-0-0); (3rd Application): 1.25 (46-0-0); Option 2: (1st Application): 0.75 (16-20-0); (2nd Application): 2.25 (46-0-0); (3rd Application): 2.00 (46-0-0); 0.25 (0-0-60); Option 3: (1st Application): 2.25 (46-0-0); 1.00 (0-18-0); (2nd Application): 1.25 (46-0-0); (3rd Application): 1.25 (46-0-0); 0.25 (0-0-60); | 110 | 7 | 7 | Option 1: (1st Application): 1.00 (14-14-14); 2.25 (46-0-0); (2nd Application): 1.25 (46-0-0); (3rd Application): 1.25 (46-0-0); Option 2: (1st Application): 0.75 (16-20-0); (2nd Application): 4.75 (46-0-0); (3rd Application): 0.25 (0-0-60); Option 3: (1st Application): 2.50 (46-0-0); 1.00 (0-18-0); (2nd Application): 1.25 (46-0-0); (3rd Application): 1.25 (46-0-0); 0.25 (0-0-60); | 120 | 7 | 7 | Option 1: (1st Application): 1.00 (14-14-14); 2.50 (46-0-0); (2nd Application): 1.50 (46-0-0); (3rd Application): 1.50 (46-0-0); Option 2: (1st Application): 0.75 (16-20-0); (2nd Application): 3.00 (46-0-0); (3rd Application): 2.00 (46-0-0); 0.25 (0-0-60); Option 3: (1st Application): 2.75 (46-0-0); 1.00 (0-18-0); (2nd Application): 1.50 (46-0-0); (3rd Application): 1.50 (46-0-0); 0.25 (0-0-60); | | | | | |
| Taboc | _no interview_ | 071 | M-L-ML-MH | 90 | 40 | 20 | Option 1: (1st Application): 3.00 (14-14-14); 1.25 (46-0-0); 2.25 (0-18-0); (2nd Application): 1.00 (46-0-0); (3rd Application): 1.00 (46-0-0); Option 2: (1st Application): 4.00 (16-20-0); (2nd Application): 1.50 (46-0-0); (3rd Application): 1.00 (46-0-0); 0.75 (0-0-60); Option 3: (1st Application): 2.00 (46-0-0); 4.50 (0-18-0); (2nd Application): 1.00 (46-0-0); (3rd Application): 1.00 (46-0-0); 0.75 (0-0-60); | 100 | 40 | 20 | Option 1: (1st Application): 3.00 (14-14-14); 1.30 (46-0-0); 2.22 (0-18-0); (2nd Application): 1.09 (46-0-0); (3rd Application): 1.09 (46-0-0); Option 2: (1st Application): 4.00 (16-20-0); (2nd Application): 2.00 (46-0-0); (3rd Application): 1.00 (46-0-0); 0.75 (0-0-60); Option 3: (1st Application): 2.25 (46-0-0); 4.50 (0-18-0); (2nd Application): 1.25 (46-0-0); (3rd Application): 1.25 (46-0-0); 0.75 (0-0-60); | 110 | 50 | 30 | Option 1: (1st Application): 4.50 (14-14-14); 1.25 (46-0-0); 2.22 (0-18-0); (2nd Application): 1.20 (46-0-0); (3rd Application): 1.20 (46-0-0); Option 2: (1st Application): 5.00 (16-20-0); (2nd Application): 2.00 (46-0-0); (3rd Application): 1.25 (46-0-0); 1.00 (0-0-60); Option 3: (1st Application): 2.50 (46-0-0); 5.75 (0-18-0); (2nd Application): 1.25 (46-0-0); (3rd Application): 1.25 (46-0-0); 1.00 (0-0-60); | 120 | 50 | 30 | Option 1: (1st Application): 4.50 (14-14-14); 1.50 (46-0-0); 2.25 (0-18-0); (2nd Application): 1.50 (46-0-0); (3rd Application): 1.50 (46-0-0); Option 2: (1st Application): 5.00 (16-20-0); (2nd Application): 2.00 (46-0-0); (3rd Application): 1.50 (46-0-0); 1.00 (0-0-60); Option 3: (1st Application): 2.75 (46-0-0); 5.25 (0-18-0); (2nd Application): 1.50 (46-0-0); (3rd Application): 1.50 (46-0-0); 1.00 (0-0-60); | | | | | |
| Taboc | Julie Baculio | 092 | M-ML-MH-H | 70 | 20 | 7 | Option 1: (1st Application): 1.00 (14-14-14); 1.25 (46-0-0); 1.50 (0-18-0); (2nd Application): 1.00 (46-0-0); (3rd Application): 1.00 (46-0-0); Option 2: (1st Application): 2.00 (16-20-0); (2nd Application): 1.50 (46-0-0); (3rd Application): 1.00 (46-0-0); 0.25 (0-0-60); Option 3: (1st Application): 1.50 (46-0-0); 2.25 (0-18-0); (2nd Application): 1.00 (46-0-0); (3rd Application): 1.00 (46-0-0); 0.25 (0-0-60); | 80 | 20 | 7 | Option 1: (1st Application): 1.00 (14-14-14); 1.50 (46-0-0); 1.50 (0-18-0); (2nd Application): 1.00 (46-0-0); (3rd Application): 1.00 (46-0-0); Option 2: (1st Application): 2.00 (16-20-0); (2nd Application): 2.00 (46-0-0); (3rd Application): 1.00 (46-0-0); 0.25 (0-0-60); Option 3: (1st Application): 1.75 (46-0-0); 2.25 (0-18-0); (2nd Application): 1.00 (46-0-0); (3rd Application): 1.00 (46-0-0); 0.25 (0-0-60); | 90 | 30 | 7 | Option 1: (1st Application): 1.00 (14-14-14); 1.75 (46-0-0); 2.75 (0-18-0); (2nd Application): 1.00 (46-0-0); (3rd Application): 1.00 (46-0-0); Option 2: (1st Application): 3.00 (16-20-0); (2nd Application): 2.00 (46-0-0); (3rd Application): 1.00 (46-0-0); 0.25 (0-0-60); Option 3: (1st Application): 2.00 (46-0-0); 3.50 (0-18-0); (2nd Application): 1.00 (46-0-0); (3rd Application): 1.00 (46-0-0); 0.25 (0-0-60); | 100 | 30 | 7 | Option 1: (1st Application): 1.00 (14-14-14); 2.00 (46-0-0); 2.75 (0-18-0); (2nd Application): 1.25 (46-0-0); (3rd Application): 1.25 (46-0-0); Option 2: (1st Application): 3.00 (16-20-0); (2nd Application): 2.00 (46-0-0); (3rd Application): 1.50 (46-0-0); 0.25 (0-0-60); Option 3: (1st Application): 2.25 (46-0-0); 3.50 (0-18-0); (2nd Application): 1.25 (46-0-0); (3rd Application): 1.25 (46-0-0); 0.25 (0-0-60); | | | | | |

FOR THE LEGEND

Fertilizer Guide for Rice

For Transplanted

after second harrowing: 10-20 bags organic fertilizer (basal)

Inorganic Fertilizer:

- 1st Application: after last harrowing
2nd Application: 10-14 DAT (days after transplanting)
3rd Application: 20-25 DAT (days after transplanting)

For Direct Seeded

after second harrowing: 10-20 bags organic fertilizer (basal)

Inorganic Fertilizer:

- 1st Application: 10 - 15 DAS (days after seeding)
2nd Application: 20-30 DAS (days after seeding)
3rd Application: 35-45 DAS (days after seeding)