Department of Agriculture
BUREAU OF SOILS AND WATER MANAGEMENT
"NATIONAL SOIL SAMPLING AND TESTING FOR FERTILITY AND CROP SUITABILITY ASSESSMENT"
FERTILIZER-GUIDE FOR RICE
PROVINCE OF ORIENTAL MINDORO

Additional 10-20 bags/ha Organic Fertilizer is recommended for all options

Municipality	Fertilizer-Guide Group Number	Description Texture-N-P-K	INBRED RICE WET SEASON Nutrient Requirement N	P₂O₅	K₂O	INBRED RICE WET SEASON Recommended Fertilizer Rate (bags/hectare)	INBRED RICE DRY SEASON Nutrient Requirement N	P₂O₅	K₂O	INBRED RICE DRY SEASON Recommended Fertilizer Rate (bags/hectare)	HYBRID RICE WET SEASON Nutrient Requirement N	P₂O₅	K₂O	HYBRID RICE WET SEASON Recommended Fertilizer Rate (bags/hectare)	HYBRID RICE DRY SEASON Nutrient Requirement N	P₂O₅	K₂O	HYBRID RICE DRY SEASON Recommended Fertilizer Rate (bags/hectare)

Due to the extreme width and structural complexity of this table, it is reproduced below with nutrient-requirement values (N, P_2O_5, K_2O) stated in LaTeX.

Row 1 — VICTORIA, Group 077, M-L-H-L

Inbred Wet Season: $N=90$, $P_2O_5=7$, $K_2O=60$
Recommended Fertilizer Rate: Option 1: (1st Application): 1.00 (14-14-14-); 1.75 (46-0-0); (2nd Application): 1.00 (46-0-0); (3rd Application): 1.00 (46-0-0); 2.00 (0-0-60); Option 2: (1st Application): 0.75 (16-20-0); (2nd Application): 2.00 (46-0-0); (3rd Application): 1.75 (46-0-0); 2.00 (0-0-60); Option 3: (1st Application): 2.00 (46-0-0); 1.00 (0-18-0); (2nd Application): 1.00 (46-0-0); (3rd Application): 1.00 (46-0-0); 2.00 (0-0-60);

Inbred Dry Season: $N=100$, $P_2O_5=7$, $K_2O=60$
Recommended Fertilizer Rate: Option 1: (1st Application): 1.00 (14-14-14-); 2.00 (46-0-0); (2nd Application): 1.25 (46-0-0); (3rd Application): 1.25 (46-0-0); 2.00 (0-0-60); Option 2: (1st Application): 0.75 (16-20-0); (2nd Application): 2.25 (46-0-0); (3rd Application): 2.00 (46-0-0); 2.00 (0-0-60); Option 3: (1st Application): 2.25 (46-0-0); 1.00 (0-18-0); (2nd Application): 1.25 (46-0-0); (3rd Application): 1.25 (46-0-0); 2.00 (0-0-60);

Hybrid Wet Season: $N=110$, $P_2O_5=7$, $K_2O=70$
Recommended Fertilizer Rate: Option 1: (1st Application): 1.00 (14-14-14-); 2.25 (46-0-0); (2nd Application): 1.25 (46-0-0); (3rd Application): 1.25 (46-0-0); 2.25 (0-0-60); Option 2: (1st Application): 0.75 (16-20-0); (2nd Application): 4.75 (46-0-0); (3rd Application): 2.50 (0-0-60); Option 3: (1st Application): 2.50 (46-0-0); 1.00 (0-18-0); (2nd Application): 1.25 (46-0-0); (3rd Application): 1.25 (46-0-0); 2.50 (0-0-60);

Hybrid Dry Season: $N=120$, $P_2O_5=7$, $K_2O=70$
Recommended Fertilizer Rate: Option 1: (1st Application): 1.00 (14-14-14-); 2.50 (46-0-0); (2nd Application): 1.50 (46-0-0); (3rd Application): 1.50 (46-0-0); 2.25 (0-0-60); Option 2: (1st Application): 0.75 (16-20-0); (2nd Application): 3.00 (46-0-0); (3rd Application): 2.00 (46-0-0); 2.50 (0-0-60); Option 3: (1st Application): 2.75 (46-0-0); 1.00 (0-18-0); (2nd Application): 1.50 (46-0-0); (3rd Application): 1.50 (46-0-0); 2.50 (0-0-60);

Row 2 — VICTORIA, Group 065, M-L-L-L

Inbred Wet Season: $N=90$, $P_2O_5=60$, $K_2O=60$
Recommended Fertilizer Rate: Option 1: (1st Application): 8.75 (14-14-14-); (2nd Application): 0.75 (46-0-0); (3rd Application): 0.75 (46-0-0); Option 2: (1st Application): 6.00 (16-20-0); (2nd Application): 1.00 (46-0-0); (3rd Application): 1.00 (46-0-0); 2.00 (0-0-60); Option 3: (1st Application): 2.00 (46-0-0); 6.75 (0-18-0); (2nd Application): 1.00 (46-0-0); (3rd Application): 1.00 (46-0-0); 2.00 (0-0-60);

Inbred Dry Season: $N=100$, $P_2O_5=60$, $K_2O=60$
Recommended Fertilizer Rate: Option 1: (1st Application): 8.75 (14-14-14-); (2nd Application): 1.00 (46-0-0); (3rd Application): 1.00 (46-0-0); Option 2: (1st Application): 6.00 (16-20-0); (2nd Application): 1.25 (46-0-0); (3rd Application): 1.00 (46-0-0); 2.00 (0-0-60); Option 3: (1st Application): 2.25 (46-0-0); 6.75 (0-18-0); (2nd Application): 1.25 (46-0-0); (3rd Application): 1.25 (46-0-0); 2.00 (0-0-60);

Hybrid Wet Season: $N=110$, $P_2O_5=70$, $K_2O=70$
Recommended Fertilizer Rate: Option 1: (1st Application): 10.00 (14-14-14-); (2nd Application): 1.00 (46-0-0); (3rd Application): 1.00 (46-0-0); Option 2: (1st Application): 7.00 (16-20-0); (2nd Application): 1.50 (46-0-0); (3rd Application): 1.00 (46-0-0); 2.50 (0-0-60); Option 3: (1st Application): 2.50 (46-0-0); 8.00 (0-18-0); (2nd Application): 1.25 (46-0-0); (3rd Application): 1.25 (46-0-0); 2.50 (0-0-60);

Hybrid Dry Season: $N=120$, $P_2O_5=70$, $K_2O=70$
Recommended Fertilizer Rate: Option 1: (1st Application): 10.00 (14-14-14-); (2nd Application): 1.25 (46-0-0); (3rd Application): 1.25 (46-0-0); Option 2: (1st Application): 7.00 (16-20-0); (2nd Application): 2.00 (46-0-0); (3rd Application): 1.00 (46-0-0); 2.50 (0-0-60); Option 3: (1st Application): 2.75 (46-0-0); 8.00 (0-18-0); (2nd Application): 1.50 (46-0-0); (3rd Application): 1.50 (46-0-0); 2.50 (0-0-60);

Row 3 — VICTORIA, Group 066, M-L-L-ML

Inbred Wet Season: $N=90$, $P_2O_5=60$, $K_2O=40$
Recommended Fertilizer Rate: Option 1: (1st Application): 5.75 (14-14-14-); 2.25 (0-18-0); (2nd Application): 1.25 (46-0-0); (3rd Application): 1.25 (46-0-0); Option 2: (1st Application): 6.00 (16-20-0); (2nd Application): 1.00 (46-0-0); (3rd Application): 1.00 (46-0-0); 1.50 (0-0-60); Option 3: (1st Application): 2.00 (46-0-0); 6.75 (0-18-0); (2nd Application): 1.00 (46-0-0); (3rd Application): 1.00 (46-0-0); 1.50 (0-0-60);

Inbred Dry Season: $N=100$, $P_2O_5=60$, $K_2O=40$
Recommended Fertilizer Rate: Option 1: (1st Application): 5.75 (14-14-14-); 2.25 (0-18-0); (2nd Application): 1.50 (46-0-0); (3rd Application): 1.50 (46-0-0); Option 2: (1st Application): 6.00 (16-20-0); (2nd Application): 1.25 (46-0-0); (3rd Application): 1.00 (46-0-0); 1.50 (0-0-60); Option 3: (1st Application): 2.25 (46-0-0); 6.75 (0-18-0); (2nd Application): 1.25 (46-0-0); (3rd Application): 1.25 (46-0-0); 1.50 (0-0-60);

Hybrid Wet Season: $N=110$, $P_2O_5=70$, $K_2O=50$
Recommended Fertilizer Rate: Option 1: (1st Application): 7.25 (14-14-14-); 2.25 (0-18-0); (2nd Application): 1.50 (46-0-0); (3rd Application): 1.50 (46-0-0); Option 2: (1st Application): 7.00 (16-20-0); (2nd Application): 1.50 (46-0-0); (3rd Application): 1.00 (46-0-0); 1.75 (0-0-60); Option 3: (1st Application): 2.50 (46-0-0); 8.00 (0-18-0); (2nd Application): 1.25 (46-0-0); (3rd Application): 1.25 (46-0-0); 1.75 (0-0-60);

Hybrid Dry Season: $N=120$, $P_2O_5=70$, $K_2O=50$
Recommended Fertilizer Rate: Option 1: (1st Application): 7.25 (14-14-14-); 2.25 (0-18-0); (2nd Application): 1.50 (46-0-0); (3rd Application): 1.50 (46-0-0); Option 2: (1st Application): 7.00 (16-20-0); (2nd Application): 2.00 (46-0-0); (3rd Application): 1.00 (46-0-0); 1.75 (0-0-60); Option 3: (1st Application): 2.75 (46-0-0); 8.00 (0-18-0); (2nd Application): 1.50 (46-0-0); (3rd Application): 1.50 (46-0-0); 1.75 (0-0-60);

Row 4 — VICTORIA, Group 073, M-L-MH-L

Inbred Wet Season: $N=90$, $P_2O_5=20$, $K_2O=60$
Recommended Fertilizer Rate: Option 1: (1st Application): 3.00 (14-14-14-); 1.25 (46-0-0); (2nd Application): 1.00 (46-0-0); (3rd Application): 1.00 (46-0-0); 1.50 (0-0-60); Option 2: (1st Application): 2.00 (16-20-0); (2nd Application): 2.25 (46-0-0); (3rd Application): 1.00 (46-0-0); 2.00 (0-0-60); Option 3: (1st Application): 2.00 (46-0-0); 2.25 (0-18-0); (2nd Application): 1.00 (46-0-0); (3rd Application): 1.00 (46-0-0); 2.00 (0-0-60);

Inbred Dry Season: $N=100$, $P_2O_5=20$, $K_2O=60$
Recommended Fertilizer Rate: Option 1: (1st Application): 3.00 (14-14-14-); 1.50 (46-0-0); (2nd Application): 1.25 (46-0-0); (3rd Application): 1.25 (46-0-0); 1.50 (0-0-60); Option 2: (1st Application): 2.00 (16-20-0); (2nd Application): 2.00 (46-0-0); (3rd Application): 1.75 (46-0-0); 2.00 (0-0-60); Option 3: (1st Application): 2.25 (46-0-0); 2.25 (0-18-0); (2nd Application): 1.25 (46-0-0); (3rd Application): 1.25 (46-0-0); 2.00 (0-0-60);

Hybrid Wet Season: $N=110$, $P_2O_5=30$, $K_2O=70$
Recommended Fertilizer Rate: Option 1: (1st Application): 4.50 (14-14-14-); 1.25 (46-0-0); (2nd Application): 1.25 (46-0-0); (3rd Application): 1.25 (46-0-0); 1.50 (0-0-60); Option 2: (1st Application): 3.00 (16-20-0); (2nd Application): 2.00 (46-0-0); (3rd Application): 1.75 (46-0-0); 2.50 (0-0-60); Option 3: (1st Application): 2.50 (46-0-0); 3.50 (0-18-0); (2nd Application): 1.25 (46-0-0); (3rd Application): 1.25 (46-0-0); 2.50 (0-0-60);

Hybrid Dry Season: $N=120$, $P_2O_5=30$, $K_2O=70$
Recommended Fertilizer Rate: Option 1: (1st Application): 4.50 (14-14-14-); 1.50 (46-0-0); (2nd Application): 1.50 (46-0-0); (3rd Application): 1.50 (46-0-0); 1.50 (0-0-60); Option 2: (1st Application): 3.00 (16-20-0); (2nd Application): 2.25 (46-0-0); (3rd Application): 2.00 (46-0-0); 2.50 (0-0-60); Option 3: (1st Application): 2.75 (46-0-0); 3.50 (0-18-0); (2nd Application): 1.50 (46-0-0); (3rd Application): 1.50 (46-0-0); 2.50 (0-0-60);

Row 5 — VICTORIA, Group 074, M-L-MH-ML

Inbred Wet Season: $N=90$, $P_2O_5=20$, $K_2O=40$
Recommended Fertilizer Rate: Option 1: (1st Application): 3.00 (14-14-14-); 1.25 (46-0-0); (2nd Application): 1.00 (46-0-0); (3rd Application): 1.00 (46-0-0); 0.75 (0-0-60); Option 2: (1st Application): 2.00 (16-20-0); (2nd Application): 2.25 (46-0-0); (3rd Application): 1.00 (46-0-0); 1.50 (0-0-60); Option 3: (1st Application): 2.00 (46-0-0); 2.25 (0-18-0); (2nd Application): 1.00 (46-0-0); (3rd Application): 1.00 (46-0-0); 1.50 (0-0-60);

Inbred Dry Season: $N=100$, $P_2O_5=20$, $K_2O=40$
Recommended Fertilizer Rate: Option 1: (1st Application): 3.00 (14-14-14-); 1.50 (46-0-0); (2nd Application): 1.25 (46-0-0); (3rd Application): 1.25 (46-0-0); 0.75 (0-0-60); Option 2: (1st Application): 2.00 (16-20-0); (2nd Application): 2.00 (46-0-0); (3rd Application): 1.75 (46-0-0); 1.50 (0-0-60); Option 3: (1st Application): 2.25 (46-0-0); 2.25 (0-18-0); (2nd Application): 1.25 (46-0-0); (3rd Application): 1.25 (46-0-0); 1.50 (0-0-60);

Hybrid Wet Season: $N=110$, $P_2O_5=30$, $K_2O=50$
Recommended Fertilizer Rate: Option 1: (1st Application): 4.50 (14-14-14-); 1.25 (46-0-0); (2nd Application): 1.25 (46-0-0); (3rd Application): 1.25 (46-0-0); 0.75 (0-0-60); Option 2: (1st Application): 3.00 (16-20-0); (2nd Application): 2.00 (46-0-0); (3rd Application): 1.75 (46-0-0); 1.75 (0-0-60); Option 3: (1st Application): 2.50 (46-0-0); 3.50 (0-18-0); (2nd Application): 1.25 (46-0-0); (3rd Application): 1.25 (46-0-0); 1.75 (0-0-60);

Hybrid Dry Season: $N=120$, $P_2O_5=30$, $K_2O=50$
Recommended Fertilizer Rate: Option 1: (1st Application): 4.50 (14-14-14-); 1.50 (46-0-0); (2nd Application): 1.25 (46-0-0); (3rd Application): 1.50 (46-0-0); 0.75 (0-0-60); Option 2: (1st Application): 3.00 (16-20-0); (2nd Application): 2.25 (46-0-0); (3rd Application): 2.00 (46-0-0); 1.75 (0-0-60); Option 3: (1st Application): 2.75 (46-0-0); 3.50 (0-18-0); (2nd Application): 1.50 (46-0-0); (3rd Application): 1.50 (46-0-0); 1.75 (0-0-60);