
### LAND SUITABILITY MAP

### **ABACA**

## LAND RESOURCES EVALUATION AND SUITABILITY ASSESSMENT OF STRATEGIC PRODUCTION AREAS


### **ZAMBOANGA CITY**





### MAP INDEX

# LAND RESOURCES EVALUATION AND SUITABILITY ASSESSMENT OF STRATEGIC PRODUCTION AREAS ZAMBOANGA CITY



## LAND SUITABILITY MAP FOR **ABACA**

### LAND RESOURCES EVALUATION AND SUITABILITY ASSESSMENT OF STRATEGIC PRODUCTION AREAS ZAMBOANGA CITY, REGION IX

#### **EXTENT OF SUITABILITY FOR ABACA PRODUCTION BY MUNICIPALITY**

					·	EX	PANSION	AREA (H	a)		•	CONFLIC'	T RESOL	UTION A	REA (Ha)		TOTAL
MUNICIPALITY	EXIST	ING ABAO	CA (Ha)	TOTAL EXISTING AREA (Ha)	Coco	nut	Shrub unman	,	Grassl unmana	·	Co	rn	•	y rice, rigated	Other	crops	POTENTIAL EXPANSION AREA (Ha)
	<b>S1</b>	<b>S2</b>	<b>S</b> 3	1	<b>S1</b>	<b>S2</b>	<b>S1</b>	<b>S2</b>	S1	<b>S2</b>	S1	<b>S2</b>	<b>S1</b>	<b>S2</b>	<b>S1</b>	<b>S2</b>	- АКЕА (Па)
ZAMBOANGA CITY	-	_	_	-	6,382	16,241	1,148	3,722	2,686	9,772	2,487	1,378	-	-	-	8	43,823
TOTAL	-	_	-	-	6,382	16,241	1,148	3,722	2,686	9,772	2,487	1,378	-	-	-	8	43,823

Note: Delivery of abaca planting materials must be started on the onset of rainy season.

- undulating to rolling

#### AGRONOMIC REQUIREMENT OF ABACA PRODUCTION

UTILIZATI TYPE	ON SUITABILITY RATING	SLOPE (%)	SOIL DEPTH (cm)	SOIL TEXTURE	SOIL DRAINAGE	SUIL REACTION (pH)	INHERENT FERTILITY	FLOODING CLASS	EROSION CLASS	ROCK OUTCROPS	ELEVATION (masl)	ANNUAL RAINFALL (mm)	CLIMATIC TYPE
	S1	<8	>50	CL, SiCL, SCL, SC, SiC, C, HC	WD,MWD, SPD	5.6 -7.2	high	none-slight	none-slight	none-few	<500	2001-4500	II, III, IV
Abaca	S2	8 - 30	30 - 50	FSL, L, SiL, SL	PD,VPD	5.1 - 5.5 7.3 - 7.8	medium	moderate	moderate	common	500-1500	1000-2000	I, II
	S3	>30	< 30	S, LS, CSL	ED	<5.0 - > 7.9	low	severe	severe	many	>1500	<1000 >4500	
SLOPE (%) SOIL DRAINAGE			SOIL REACTION (pH)			SOIL TEXTURE							
0 - 3	level to gently slopin	ng	ED - 6	excessively drained		< 4.5 - ext	remely acid		Coarse			Fine	
3 - 8	gently sloping to und	dulating	WD - v	well drained		4.5 - 5.0 - ver	y strongly acid		S - s	and		SC - s	sandy clay

5.1 - 5.5 - strongly acid

18 - 30 - somewhat poorly drained 5.6 - 6.0 - medium acid - rolling to moderately steep - coarse sandy loam 30 - 50 poorly drained 6.1 - 6.5 - slightly acid - sandy loam > 50 VPD very poorly drained 6.6 - 7.2 - neutral very steep 7.3 - 7.8 - mildly alkaline - fine sandy loam SOIL DEPTH (cm) **SURFACE IMPEDIMENT** - moderately alkaline - loam very shallow ROCK OUTCROPS - strongly alkaline - silt loam 30 - 50 - none - few - clay loam - moderately deep - silty clay loam 10 - 30% - common - deep to very deep > 30% - sandy clay loam

- moderately well drained

#### LAND LIMITATIONS DESCRIPTION AND COMBINATIONS

ELEVATION	SOIL DRAINAGE	SOIL DEPTH	<b>SOIL EROSION</b>
El2 - 500 - 1000m or 2000 - 2500m	D2 - Somewhat poorly drained to poorly drained	Sh2 - Shallow to moderately deep (30 - 100cm)	E2 - Moderate erosion
El3 $-<500 \text{m or} > 2500 \text{m}$	D3 - Very poorly drained or excessively drained	Sh3 - Very shallow (< 30cm)	E3 - Severe erosion
SLOPE/TOPOGRAPHY	SOIL TEXTURE	ROCK OUTCROPS	FLOODING
T2 - Undulating to moderately steep	Tc - Coarse texture	Rc2 - Common	F2 - Moderate seasonal flooding
T3 - Steep to very steep		Rc3 - Many	F3 - Severe seasonal flooding

CODE	LIMITATION	CODE	LIMITATION	CODE	LIMITATION	CODE	LIMITATION
1	E2-Rc3	11	T2-E3-Sh2-Rc3	21	Т3	31	Т3-Е3
2	F2-D2	12	T2-El2-E3-Rc3	22	Т3-Е3	32	T3-E3-Rc3
3	F3-D2	13	T2-El2-E3-Sh2-Rc2	23	T3-E3-Rc2	33	T3-E3-Sh3-Rc3
4	Sh2-Rc2	14	T2-El2-E3-Sh2-Rc3	24	T3-E3-Sh2-Rc3	34	T3-El2-E3-Sh3-Rc3
5	T2	15	T2-El2-Sh2-Rc2	25	T3-E3-Sh3-Rc2	35	T3-El3
6	T2-E2-Sh2-Rc2	16	T2-El2-Sh2-Rc3	26	T3-E3-Sh3-Rc3	36	Tc
7	T2-E3	17	T2-F2-D2	27	T3-El2-E3-Sh3-Rc2		
8	T2-E3-Rc2	18	T2-F3-D2	28	T3-El2-E3-Sh3-Rc3		
9	T2-E3-Rc3	19	T2-Sh2-Rc2	29	T3-F3-D2		
<i>10</i>	T2-E3-Sh2-Rc2	20	T2-Sh2-Rc3	30	Т3		

CODE	LANDUSE
4	Corn
82	Cacao
105	Fruit trees, mixed
116	Coconut
126	Grassland
134	Shrubs, unmanaged
137	Rubber

loamy sand

silty clay

heavy clay

- clay

#### **SUITABILITY CLASSES:**

Highly Suitable (S1) Land having no significant limitation to sustained application of a given use, or only minor limitations that will not significantly reduce productivity or benefits and will not raise inputs above an acceptable level.

Marginally Suitable (S3) Land having limitations which in aggregate are severe for sustained application of a given use and will so reduce productivity or benefits, or increase required inputs, that this expenditure will be only marginally justified.

**Moderately Suitable (S2)** Land having limitation which in aggregate are moderately severe for sustained application of a given use; the limitation will reduce productivity or benefits and increase required inputs to the extent that the overall advantage to be gained from the use, although still attractive, will be appreciably inferior to that expected on class S1 land.

**Not Suitable / Not Relevant** Land having limitations which may be surmountable in time but which cannot be corrected with existing knowledge at currently acceptable cost; the limitations are so severe as to preclude successful sustained use of the land in the given manner. Existing forest, shrubland greater than 18% slope, irrigated paddy rice and miscellaneous land types such as built up areas, roads, etc are considered as not relevant.

#### **CLIMATE TYPE**


TYPE I: Two pronouced season, dry from November to April and TYPE II: No dry season with a very pronounced maximum rain wet during the rest of the year. Maximum rain period is from June to September

period from December to February. There is not a single dry month. Maximum monthly rainfall occurs during the period from March to May.

**TYPE III:** No very pronounced maximum rain period, with a dry season lasting only from one to three months, either during the period from December to February or from March to May. This type resembles Type I since it has a short dry season.

**TYPE IV**: Rainfall is more or less evenly distributed throughout the year. This type resembles Type II since it has no dry

Whole part of Zamboanga City is classified as climatic Type III.



Source: PAGASA 2018, Climatological Normals (Rainfall), Philippine Atmospheric, Geophysical and Astronomical Services Administration (PAGASA), accessed 27 July 2018, <a href="https://www1.pagasa.dost.gov.ph/index.php/climate/climatological-normals">https://www1.pagasa.dost.gov.ph/index.php/climate/climatological-normals</a>.

<sup>\*</sup>establishment of shade trees prior to planting of abaca.

